# Uniqueness of Steiner Laws on Cubic Curves

*R. Padmanabhan*[*]
Department of Mathematics
University of Manitoba
Winnipeg, Manitoba R3T 2N2
Canada
E-mail: padman@cc.umanitoba.ca

*W. McCune*[†]
Mathematics and Computer Science Division
Argonne National Laboratory
Argonne, Illinois 60439-4844
U.S.A.
E-mail: mccune@mcs.anl.gov

August 6, 1998

**Abstract.** In this paper we use the Cayley-Bacharach theorem of classical algebraic geometry to construct several universal algebras on algebraic curves using divisors and complete intersection cycles and study the equational identities valid for these synthetic constructions. These results are not necessarily new, and in fact, all of them may be "easily" provable by resorting to such powerful tools as the Riemann-Roch theorem, the $\mathcal{P}$-function of Weierstrass, the rigidity lemma, Euler numbers, Lefschetz fixed-point theorem and so on. However, our equational proofs employ automated reasoning by transforming the Cayley-Bacharach Theorem into a formal implication. Beside being elementary, this approach provides new examples for model theorists and computer scientists designing theorem-provers and gives new insights and interpretations for these various geometric constructions.

## 1 Introduction

Let $C$ be a non-singular cubic curve in the complex projective plane. If $a$ and $b$ are two distinct points on the curve, let $c = a * b$ be the (unique) third point of intersection with $C$ of the line $L(a, b)$ joining $a$ and $b$. If $b = a$, then we naturally take the line $L(a, b)$ to be the tangent at $a$. More formally, the set $\{a, b, a * b\}$ is the complete intersection cycle of the curve $C$ with $L(a, b)$ counting multiplicities. If the ground field $k$ is different from complex numbers, then we insist that the points

[*]Supported by an operating grant from NSERC of Canada (OGP8215).

[†]Supported by the Mathematical, Information, and Computational Sciences Division subprogram of the Office of Computational and Technology Research, U.S. Department of Energy, under Contract W-31-109-Eng-38.

$a$ and $b$ are $k$-rational points. In that case, the unique third point a*b is obviously $k$-rational and it is this fact that makes the cubic curves very interesting. In fact, if $e$ is chosen to be an inflection point (again, $k$-rational if $k \neq C$), then the composite term $(a * b) * e$ is the classical Poincare group law on $C$. It is clear that the rational binary operation "$*$" viewed as a mapping from $C \times C$ to $C$ (or from $C(k) \times C(k)$ to $C(k)$ if $k$ is not algebraically closed) satisfies the following universal identities: $a * b = b * a$, $a * (b * a) = b$. This is an example of a binary Steiner law and the idempotents of $*$ are precisely the inflection points of the curve (see Fig. 1). More generally, an $n$-ary Steiner law $f(x_1, x_2, \ldots, x_n)$ on a projective curve $C$ over $k$ is a totally symmetric rational $n$-ary function $f$ from $C^n$ to $C$ satisfying the universal identity

$$f(x_1, x_2, ..., x_{n-1}, f(x_1, x_2, ..., x_n)) = x_n.$$

An element $e$ in $C$ is called an idempotent for $f$ if $f(e, e, ...e) = e$. In this paper, we prove that if $f$ and $g$ are two $n$-ary Steiner laws on an elliptic curve $C$ sharing a common idempotent then $f = g$. First, we extract a special case of the inference rule (gL) — indeed, a fragment of the powerful rigidity lemma — from the Cayley-Bacharach Theorem of the classical algebraic geometry. This rule is implemented in OTTER, a first-order theorem-proving program. Then we use OTTER to automate the proofs of the uniqueness of the 5-ary Steiner laws definable on an elliptic curves. Very much like the binary case, this theorem provides algebraic characterizations of synthetic geometric constructions involving the intersection cycles of cubics with algebraic curves of higher degrees. The well-known theorem of the uniqueness of the group law on such a curve is an extreme special case of this result.

A set $P$ of $p$ points in $PG(2, k)$ — a projective plane over a field $k$, is said to have the Cayley-Bacharach property (CB-property) of degree $d$ if any plane curve of degree $d$ passing through all but one point of $P$ necessarily contains the whole of $P$. The Cayley-Bacharach Theorem of algebraic geometry (see e.g. [1, 2]) says that if $P$ is a set of $mn$ points which is a complete intersection cycle of two curves of degrees $m$ and $n$, then $P$ has the CB-property of degree $m + n - 3$. This classical result is rife with rich rational universal algebras (i.e., rational constructions that yield unique points). In this paper, we employ the techniques of Bezout theorem and the CB-theorem to prove that every algebraic curve induces a rational Steiner operation on cubic curves via a complete intersection cycle (see e.g. Fig. 1 for the binary linear process and Fig. 4 for the 5-ary conic process).

All the proofs in this paper reside completely within the framework of first order logic with equality. First we show that the Cayley-Bacharach Theorem implies the validity of a formal implication which is a fragment of the rule (gL) or the "Term Condition" — an algebraic avatar of the rigidity lemma of algebraic geometry. Then we exploit this in finding interrelations among various algebraic laws of different arities obtained via the Cayley-Bacharach constructions on algebraic curves. Normally, one would employ the parameters of the elliptic functions of Weierstrass, the group law via the Riemann-Roch theorem or the so-called AF+BG theorem of Max Noether etc., to prove such results in the projective geometry over elliptic curves.




Figure 1: Chord-Tangent Construction

## 2 Cayley-Bacharach Theorem Implies the $=\!(gL)\!\Rightarrow$ Rule for Cubics

**Cayley-Bacharach Theorem**. *If $P$ is a set of $mn$ points which is a complete intersection cycle of two curves of degrees $m$ and $n$, then any plane curve of degree $m+n+3$ passing through all but one point of $P$ necessarily contains the whole of $P$.*

**Theorem 1**. *Let $C$ be a nonsingular cubic curve over the complex projective plane and let "$*$" be the binary morphism of chord-tangent construction. Then the algebra $\langle C; * \rangle$ satisfies the implication*

$$(a*b)*c = (a*d)*e \quad \Rightarrow \quad (x*b)*c = (x*d)*e.$$

*Proof.* Let $Q_1$ be the quartic formed by the four lines $\{1 \cup 2 \cup 3 \cup 4\}$, let $Q_2$ be the quartic formed by the four lines $\{5 \cup 6 \cup 7 \cup 8\}$, and let $C$ be the given nonsingular cubic curve. See Fig. 2. We have

$$\begin{aligned} C \cap Q_1 &= \{a, d, a*d, c, a*b, (a*b)*c, e, x*d, (x*d)*e, x, b, x*b\} \\ C \cap Q_2 &= \{e, a*d, (a*d)*e, x, d, x*d, a, b, a*b, c, x*b, (x*b)*c\} \end{aligned}$$

Hence, if $(a*b)*c = (a*d)*e$, then both $Q_1$ and $Q_2$ share 11 common points with the base cubic $C$. Here both $Q_1$ and $Q_2$ are quartics and so, by the Cayle-Bacharach Theorem they must share the 12th common point as well. Thus $(x*b)*c = (x*d)*e$. This completes the proof of the implication

$$(a*b)*c = (a*d)*e \quad \Rightarrow \quad (x*b)*c = (x*d)*e.$$

In what follows, we call this implication the "basic (gL) rule for cubics" or simply "basic (gL)". This is only a special case of the full version of the rule (gL) which, in turn, is a modern avatar of the powerful rigidity lemma of complete varieties:




Figure 2: Basic (gL)

$f(x,b) = c \Rightarrow f(x,z) = f(y,z)$ for all terms $f$ in the mathematical structure in question (see [7], page 37 and [11] for more details and references about the related conditions like the term condition).

**Theorem 2**. *Basic (gL) for cubics* $\Rightarrow$ *the identity* $((u*v)*w)*t = ((t*v)*w)*u$.

*Proof.* (See the Appendix for an OTTER proof.) Thanks to the commutative and Steiner law, we have

$$(w*((t*v)*w))*t = (w*((u*v)*w))*u$$

because both sides reduce to the variable $v$. This equality looks like the left hand side of the implication basic (gL) with the following identifications:

$$a = w, b = (t*v)*w, c = t, d = (u*v)*w, e = u.$$

Hence, by the conclusion of the implication basic (gL), we have

$$(x*b)*c = (x*d)*e$$

for all points $x$ on the cubic. In other words, we have

$$(x*((t*v)*w))*t = (x*((u*v)*w))*u.$$

Substituting $x = t$ and simplifying, we get

$$(t*v)*w = (t*((u*v)*w))*u$$

which, modulo the commutative and Steiner laws, is the same as

$$((t * v) * w) * u = ((u * v) * w) * t.$$

**Theorem 3**. *The binary Steiner law $*$ on a non-singular cubic curve satisfies the medial identity $(x * y) * (z * u) = (x * z) * (y * u)$.* (See Fig. 3.)




Figure 3: The Medial Identity

*Proof.* (See the Appendix for an OTTER proof.) It is enough to derive the medial law from the basic (gL). We work backward to show how naturally the implication of basic (gL) works to prove such consequences. We want to derive the identity $(x * y) * (z * u) = (x * z) * (y * u)$. This equality looks exactly like the conclusion of basic (gL) and hence all we need to do is to find a suitable term $a = a(y, z, u)$ such that the relation $(a * y) * (z * u) = (a * z) * (y * u)$. The term $a = y * z$ would do the job because in this case both sides reduce to $u$.

*Historical Remark.* The validity of the median law for $*$ was first proved for plane cubics by Etherington [2]. See Knapp [4] for a different and a rather complete proof including the cases where two or more points may coincide. The first automated proof of the median law using the rule (gL) was given in [11] and [12].

To further demonstrate the impact of the Cayley-Bacharach Theorem on the geometry of the cubic curves, we show that the formal property of basic (gL) for a binary Steiner law $*$ along with a mild group-like homomorphism connection with $+$, a binary law of composition does characterize the $+$ as the (unique) group law. More precisely, we prove the following

**Theorem 4**. *Let $\Sigma$ be*

$$\{x * (y * x) = y, x * y = y * x, e * e = e, (x * e) + (e * y) = x * y\}$$

*Then $\Sigma \Rightarrow_{(basic-gL)} x + y = e * (x * y)$, where $+$ is an abelian group law.*

*Proof* (found by Otter 3.0.5b on at 1.94 seconds; see Sec. 3 for an explanation of the proof notation).

| | | |
|---|---|---|
| 3 | $x*(y*x)=y$ | |
| 5 | $x*y=y*x$ | |
| 6 | $e*e=e$ | |
| 8 | $(x*e)+(e*y)=x*y$ | |
| 10 | $(x*y)*z=(x*u)*v \;\rightarrow\; (w*y)*z=(w*u)*v$ | |
| 11 | $(x*y)*x=y$ | $[3 \rightarrow 3]$ |
| 13 | $(x*y)*y=x$ | $[3 \rightarrow 5$, flip$]$ |
| 16,15 | $x*(x*y)=y$ | $[5 \rightarrow 3]$ |
| 18,17 | $x+(e*y)=(x*e)*y$ | $[13 \rightarrow 8]$ |
| 19 | $(x*e)*y=(e*x)*y$ | $[11 \rightarrow 8 : 18]$ |
| 26 | $(e*x)*y=(x*e)*y$ | [flip 19] |
| 27 | $x+y=(x*e)*(e*y)$ | $[15 \rightarrow 17]$ |
| 44 | $(x*e)*(y*(e*x))=y$ | $[3 \rightarrow 26$, flip$]$ |
| 50 | $(x*((y*z)*u))*y=(x*z)*u$ | [10,11] |
| 296 | $(x*(e*y))*e=(x*e)*y$ | $[6 \rightarrow 50]$ |
| 303 | $(x*y)*z=(x*e)*(y*(e*z))$ | $[44 \rightarrow 50]$ |
| 349 | $(x*e)*(y*(e*z))=(x*y)*z$ | [flip 303] |
| 424,423 | $(x*e)*y=e*(x*(e*y))$ | $[5 \rightarrow 296$, flip$]$ |
| 445,444 | $(x*y)*z=e*(x*(e*(y*(e*z))))$ | $[349 : 424$, flip$]$ |
| 492 | $x+y=e*(x*y)$ | $[27 : 445,16,16]$ |

To complete the proof that + defines a group law, one notices that the associativity is simply the identity of Theorem 2:

$$\begin{aligned} \text{x + (y + z)} &= \text{e * (x * (e * (y * z)))} \\ &= \text{e * (z * (e * (y * x)))} \\ &= \text{z + (y + x)} \end{aligned}$$

and hence the binary operation + is both associative and commutative. It is clear that $x+e=x$. Finally, defining $x'$ as $e*x$, we have $x+x'=x+(e*x)=e*(x*(e*x))=e*e=e$.

*Remark.* Contrast this with the results in the Appendix.

## 3 OTTER and the Implementation of the Rule (gL)

Let us now compare the basic (gL) rule with the rigidity lemma:

| | |
|---|---|
| Basic (gL) for cubics: | $(a*b)*c=(a*d)*e \Rightarrow (x*b)*c=(x*d)*e.$ |
| Full (gL) for cubics: | $F(a,b)=F(a,c) \Rightarrow F(x,b)=F(x,c)$ for all morphisms $F$ of the curve. |
| Rigidity for cubics: | $F(x,b)=c \Rightarrow F(x,z)=F(y,z)$ for all morphisms $F$ of the curve. |

Thus it is clear that while the basic (gL) for cubics deals only with the single binary operation $*$, the full (gL) — equivalent to the rigidity lemma, see Theorem 3.3 of [7] — involves all possible morphisms and thus provides the necessary

glue that binds together these various rational morphisms and gives new and elementary equational proofs to show that all these synthetic operations of higher arities can be obtained by ruler constructions. Unlike the full (gL), the basic (gL) for $*$ is of little use if we assume no further connection between $*$ and other operations (see, for example, the Appendix). Since we discuss the uniqueness of Steiner laws on cubics, we employ the full (gL) as an inference rule as well as a rewrite rule.

OTTER [6] is a computer program that attempts to prove theorems stated in first-order logic with equality. Here we restrict our attention to its capabilities in equational logic. The user inputs axioms and the denial of the goal(s), and OTTER searches for a contradiction by working both forward from the axioms and backward from the goal(s). Equational reasoning is accomplished by paramodulation and demodulation. Paramodulation is an equality substitution rule extended with unification: if the two terms in question can be made identical by instantiating variables, then equality substitution is applied to the corresponding instances. Demodulation is the use of equalities as rewrite rules to simplify other equalities. The following example illustrates the interplay between paramodulation and demodulation. Consider $\{f(x, f(g(x), y)) = y, f(u, g(u)) = e, f(w, e) = w\}$, where $e$ is a constant; OTTER can infer $x = g(g(x))$ "in one step" by unifying $f(u, g(u))$ and $f(g(x), y))$ (which instantiates $u$ to $g(x)$ and $y$ to $g(g(x))$), replacing $f(g(x), g(g(x)))$ with $e$, and then demodulating with $f(w, e) = w$.

The full rule (gL) was implemented in OTTER in two ways that are analogous to paramodulation and demodulation. Let $F[a_1, x]$ represent a term that contains a subterm $a_1$ at a particular position, with $x$ representing everything else in the term. Suppose we have $F[a_1, x] = F[a_2, y]$, (i.e., $a_1$, and $a_2$ are in corresponding positions), with $a_1$ and $a_2$ unifiable. By (gL) we infer $F[z, x'] = F[z, y']$, where $z$ is a new variable, and $x'$ and $y'$ are the appropriate instances of $x$ and $y$. For example, from

$$f(f(x, y), f(z, f(x, z))) = f(u, f(y, u)),$$

we can (gL)-infer

$$f(f(x, y), f(z, w)) = f(f(x, z), f(y, w))$$

by unifying $u$ and $f(x, z)$ and introducing the variable $w$. We also use (gL) as a rewrite rule whenever possible. That is, we rewrite $F[a, x] = F[a, y]$ to $F[z, x] = F[z, y]$ (again, $z$ is a new variable).

OTTER **Proof Notation.** Each derived clause has a justification. The notation "$m \to n$" indicates paramodulation from $m$ into $n$; ": $i, j, k, \ldots$" indicates rewriting with the equations $i, j, k, \ldots$; and ",flip" indicates that equality was reversed (usually so that the complex side occurs on the left). The justification "[(gL)" indicates the use of $=\!(gL)\!\Rightarrow$ as an inference rule, and ":(gL)" indicates its use as a rewrite rule.

## 4 Uniqueness of n-ary Steiner Law on Cubics

In this section, we show that a non-singular cubic curve admits exactly one $n$-ary Steiner law for every $n$ congruent to $2(mod\ 3)$ If $n = 2$, this is this is the usual chord-tangent process. We give a complete proof for the next case $n = 5$

(the "conic process"). The proof of the general case is very similar and can be proved by induction. Let $C$ be a non-singular cubic and let $x, y, z, t, u$ be five points on the curve. Let $Q$ be the unique conic determined by these 5 points. By the celebrated Bezout theorem of classical geometry, we have $|C \cap Q| = 6$, counting multiplicities. Let now $f(x, y, z, t, u)$ be the 5-ary morphism on $C$ defined by the complete intersection cycle

$$C \cap Q = \{x, y, z, t, u, f(x, y, z, t, u)\}.$$

Then the unique sixth point $f(x, y, z, t, u)$ can be found by a simple ruler construction as shown in the diagram below (Fig. 4). A formal proof using the rigidity lemma was given by N. S. Mendelsohn, R. Padmanabhan and B. Wolk in [8]. Here we would like to characterize the above synthetic geometric process by means of equational identities.

The 5-ary law is totally symmetric in all of its 5 arguments and every inflection point is an idempotent for $f$: $f(e, e, e, e, e) = e$. The geometric reason for this is that the intersection multiplicity at in inflection point $e$ is six. Moreover, it satisfies the Steiner identity $f(e, e, e, x, f(e, e, e, x, y)) = y$. We claim that a non-singular cubic curve over an algebraically closed field admits exactly one such 5-ary morphism.

**Lemma 1.**

$$\left\{ \begin{array}{l} f(x,y,z,u,v) = f(x,y,z,v,u) \\ f(e,e,e,e,e) = e \\ f(e,e,e,x,f(e,e,e,x,y)) = y \end{array} \right\} \Rightarrow_{(gL)} \{f(u,v,w,x,f(u,v,w,x,y)) = y\}.$$

*Proof* (found by Otter 3.0.5b on soot.mcs.anl.gov at 0.48 seconds, with a specialized search strategy).

| | | |
|---|---|---|
| 3 | $f(x,y,z,u,v) = f(x,y,z,v,u)$ | |
| 4 | $f(e,e,e,e,e) = e$ | |
| 7,6 | $f(e,e,e,x,f(e,e,e,x,y)) = y$ | |
| 9,8 | $f(e,e,e,x,f(e,e,e,y,x)) = y$ | $[3 \to 6]$ |
| 12 | $f(x,y,z,u,f(e,e,e,u,e)) = f(x,y,z,e,e)$ | $[6 \to 4$ :(gL) :(gL) :(gL), flip$]$ |
| 21 | $f(e,e,e,x,f(y,z,u,f(v,w,v_6,e,e),x)) = f(v,w,v_6,v_7,f(y,z,u,v_7,e))$ | $[8 \to 12$ :(gL) :(gL) :(gL), flip$]$ |
| 485 | $f(x,y,z,f(x,y,z,e,e),u) = f(e,e,e,e,u)$ | [(gL) 21, flip] |
| 533 | $f(x,y,z,u,f(x,y,z,u,e)) = e$ | [485 $\to$ 21 :9, flip] |
| 613 | $f(u,v,w,x,f(u,v,w,x,y)) = y$ | $[6 \to 533$ :(gL) :7] |

**Lemma 2.**

$$\left\{ \begin{array}{l} f(x,y,z,u,v) = f(x,y,z,v,u) \\ g(x,y,z,u,v) = g(x,y,z,v,u) \\ g(u,v,w,x,g(u,v,w,x,y)) = y \\ g(e,e,e,e,e) = e \end{array} \right\} \Rightarrow_{(gL)} \Rightarrow$$

$$\{f(x,y,z,u,g(v,w,v_6,u,v_7)) = f(x,y,z,v_8,g(v,w,v_6,v_8,v_7))\}$$

*Proof* (found by Otter 3.0.5b on soot.mcs.anl.gov at 0.85 seconds, with a specialized search strategy).

3 $f(x,y,z,u,v) = f(x,y,z,v,u)$

5 $g(x,y,z,u,v) = g(x,y,z,v,u)$

6 $g(e,e,e,e,e) = e$

7 $g(u,v,w,x,g(u,v,w,x,y)) = y$

9 $f(x,y,z,u,g(e,e,e,e,e)) = f(x,y,z,e,u)$ [6 → 3]

10 $f(x,y,z,e,g(u,v,w,v_6,g(u,v,w,v_6,v_7))) = f(x,y,z,v_7,g(e,e,e,e,e))$ [7 → 9, flip]

13 $f(x,y,z,e,g(u,v,w,v_6,g(u,v,w,v_7,v_6))) = f(x,y,z,v_7,g(e,e,e,e,e))$ [5 → 10]

17 $f(x,y,z,e,g(u,v,w,v_6,g(u,v,w,e,v_7))) = f(x,y,z,v_7,g(e,e,e,v_6,e))$ [(gL) 10]

33 $f(x,y,z,e,g(u,v,w,e,v_6)) = f(x,y,z,v_7,g(u,v,w,v_7,v_6))$ [13 → 17 :(gL) :(gL) :(gL) :(gL)]

40 $f(x,y,z,u,g(v,w,v_6,u,v_7)) = f(x,y,z,v_8,g(v,w,v_6,v_8,v_7))$ [33 → 33]

**Theorem 5**. *Let $S$ be the set of identities of type (5,5,0) defined by*

$$S = \left\{ \begin{array}{llll} f(e,e,e,e,e) = e, & f \text{ is symmetric}, & f(e,e,e,x,f(e,e,e,x,y)) = y, \\ g(e,e,e,e,e) = e, & g \text{ is symmetric}, & g(e,e,e,x,g(e,e,e,x,y)) = y \end{array} \right\}.$$

*Then $S$ $\Rightarrow(gL)\Rightarrow$ $\{f(x,y,z,u,v) = g(x,y,z,u,v)\}$.*

By Lemmas 1 and 2, we assume

$$f(u,v,w,x,f(u,v,w,x,y)) = y,$$
$$g(u,v,w,x,g(u,v,w,x,y)) = y,$$
$$f(x,y,z,u,g(v,w,v_6,u,v_7)) = f(x,y,z,v_8,g(v,w,v_6,v_8,v_7)).$$

*Proof* (found by Otter 3.0.5b on soot.mcs.anl.gov at 0.34 seconds, with a specialized search strategy). Full symmetry of the operations causes an explosion in the OTTER search space; to constrain the search, we incompletely specify symmetry with deduction rule 2 below.

2 $g(x,y,z,u,v) = f(x,y,z,u,v) \rightarrow g(y,z,u,v,x) = f(y,z,u,v,x)$

3 $f(e,e,e,e,e) = e$

4 $f(u,v,w,x,f(u,v,w,x,y)) = y$

5 $g(e,e,e,e,e) = e$

6 $g(u,v,w,x,g(u,v,w,x,y)) = y$

7 $f(x,y,z,u,g(v,w,v_6,u,v_7)) = f(x,y,z,v_8,g(v,w,v_6,v_8,v_7))$

10 $f(e,e,e,e,g(e,e,e,e,e)) = e$ [5 → 3]

11 $f(e,e,e,x,g(e,e,e,x,e)) = e$ [7 → 10]

12 $g(e,e,e,x,e) = f(e,e,e,x,e)$ [11 → 4, flip]

13 $g(e,e,x,e,e) = f(e,e,x,e,e)$ [12,2]

15 $f(e,e,x,e,g(e,e,x,e,e)) = e$ [13 → 4]

18 $f(e,e,x,y,g(e,e,x,y,e)) = e$ [7 → 15]

19 $g(e,e,x,y,e) = f(e,e,x,y,e)$ [18 → 4, flip]

20 $g(e,x,y,e,e) = f(e,x,y,e,e)$ [19,2]

22 $f(e,x,y,e,g(e,x,y,e,e)) = e$ [20 → 4]

25 $f(e,x,y,z,g(e,x,y,z,e)) = e$ [7 → 22]

26 $g(e,x,y,z,e) = f(e,x,y,z,e)$ [25 → 4, flip]

27 $g(x,y,z,e,e) = f(x,y,z,e,e)$ [26,2]

29 $f(x,y,z,e,g(x,y,z,e,e)) = e$ [27 → 4]

32 $f(x,y,z,u,g(x,y,z,u,e)) = e$ [7 → 29]

33 $g(x,y,z,u,e) = f(x,y,z,u,e)$ [32 → 4, flip]

34 $g(x,y,z,e,u) = f(x,y,z,e,u)$ [33,2]

| | | |
|---|---|---|
| 36 | $f(x,y,z,e,g(x,y,z,e,u)) = u$ | $[6 \to 34, \text{flip}]$ |
| 39 | $f(x,y,z,u,g(x,y,z,u,v)) = v$ | $[7 \to 36]$ |
| 40 | $g(x,y,z,u,v) = f(x,y,z,u,v)$ | $[6 \to 39, \text{flip}]$ |

Line 40 completes the proof of Theorem 5.

**Corollary 1** *$f(x,u,z,t,u) = ((x*y)*(z*t))*u$, where "$*$" stands for the binary morphism of secant-tangent construction on the cubic.*

*Proof.* Define $g(x,y,z,t,u) = ((x*y)*(z*t))*u$. It is clear that $g$ is totally symmetric and that every inflection point is an idempotent for $g$. Moreover, $g$ satisfies the 5-ary Steiner law $g(e,e,e,x,g(e,e,e,x,y)) = y$. Hence, by Theorem 2, $f = g$.

This gives the well-known ruler construction to locate the unique sixth point $f(x,y,z,t,u)$ on the cubic. See Fig. 4.




Figure 4: The Sixth Point

Thus, in particular, a non-singular cubic curve admits exactly one 5-ary totally symmetric Steiner law with flex points as its idempotents. On similar lines (or, by using a formal induction on the arity), one can easily show that for every $n \equiv 2(mod\ 3)$, a non-singular cubic curve admits exactly one totally symmetric $n$-ary Steiner law with flex points as its idempotents. Moreover, they all can be achieved by linear constructions, thanks again to the Cayley-Bacharach Theorem.

Using similar constructions, one can show that for every $n \equiv 2(mod\ 3)$, a non-singular cubic curve admits exactly one totally symmetric n-ary Steiner law with flex points as its idempotents. Indeed, such a Steiner law would be cut on a non-singular cubic by an algebraic curve of degree $d = (n+1)/3$. Once again, the fact that these operations are well-defined follows immediately thanks to the Cayley-Bacharach Theorem. Let us quickly illustrate this for, say $n = 11$, the first non-trivial case where the Cayley-Bacharach theorem really applies. If one takes a set of 11 points of general position on a non-singular cubic curve $C$, then there exist infinitely many

quartic curves passing through these 11 points. If $Q$ is one such quartic curve, it has a 12th common point with the cubic curve since $|C \cap Q| = 12$ by the Bezout Theorem (remember, throughout this paper we are denizens of the complex projective plane). Since this set $P$ of 12 points is the complete intersection cycle of two curves of degrees 3 and 4 respectively, it enjoys the CB-property of degree $3 + 4 - 3 = 4$, meaning that every quartic passing through the initial 11 points must pass through this last point as well! Bingo, we now have a well-defined universal algebra of arity 11. It is obvious that this is a totally symmetric Steiner law on the cubic. The uniqueness and linear representation all follow easily along the previous lines. Indeed, using the full version of (gL), OTTER proved the uniqueness of 8-ary Steiner law without much difficulty. Humans can easily prove the uniqueness by induction on the arity.

# Appendix

This appendix contains OTTER proofs of Theorems 2 and 3, an example of a ternary version of basic (gL), and an example showing that the full (gL) rule is more powerful than the basic (gL) rule.

### An OTTER proof of Theorem 2

*Proof* (found by Otter 3.0.5b on soot.mcs.anl.gov at 0.58 seconds).

| | | |
|---|---|---|
| 3 | $x * (y * x) = y$ | |
| 5 | $x * y = y * x$ | |
| 6 | $(x * y) * z = (x * u) * v \ \rightarrow \ (w * y) * z = (w * u) * v$ | |
| 7 | $(x * y) * x = y$ | $[3 \rightarrow 3]$ |
| 9 | $(x * y) * y = x$ | $[3 \rightarrow 5$, flip$]$ |
| 14 | $(x * ((y * z) * u)) * y = (x * z) * u$ | [6,7] |
| 35 | $(((x * y) * z) * u) * x = (u * y) * z$ | $[5 \rightarrow 14]$ |
| 1101 | $((x * y) * z) * u = ((u * y) * z) * x$ | $[35 \rightarrow 9]$ |

### An OTTER proof of Theorem 3

*Proof* (found by Otter 3.0.5b on soot.mcs.anl.gov at 0.09 seconds).

| | | |
|---|---|---|
| 3 | $x * (y * x) = y$ | |
| 6 | $(x * y) * z = (x * u) * v \ \rightarrow \ (w * y) * z = (w * u) * v$ | |
| 7 | $(x * y) * x = y$ | $[3 \rightarrow 3]$ |
| 14 | $(x * ((y * z) * u)) * y = (x * z) * u$ | [6,7] |
| 23 | $(x * (y * z)) * (u * y) = (x * u) * z$ | $[7 \rightarrow 14]$ |
| 189 | $(x * y) * (z * u) = (x * z) * (y * u)$ | $[3 \rightarrow 23]$ |

## Ternary $=\!(basic-gL)\!\Rightarrow$

This is an OTTER proof of the associativity of the ternary Mal'cev operation using a ternary version of basic (gL).

**Theorem.**

$$\left\{\begin{array}{l} m(x,y,z)=m(x,u,v) \;\rightarrow \\ \qquad\qquad m(w,y,z)=m(w,u,v) \\ m(x,y,y)=x \\ m(x,y,z)=m(z,y,x) \end{array}\right\} \Rightarrow \{m(x,y,m(z,u,v))=m(m(x,y,z),u,v)\}.$$

*Proof* (found by Otter 3.0.5b on soot.mcs.anl.gov at 0.13 seconds).

| | | |
|---|---|---|
| 3 | $m(x,y,z)=m(x,u,v) \;\rightarrow\; m(w,y,z)=m(w,u,v)$ | |
| 4 | $m(x,y,y)=x$ | |
| 6 | $m(x,y,z)=m(z,y,x)$ | |
| 7 | $m(x,x,y)=y$ | [4 → 6, flip] |
| 9 | $m(x,y,m(y,z,u))=m(x,z,u)$ | [7,3] |
| 11 | $m(x,y,m(z,u,y))=m(x,u,z)$ | [6 → 9] |
| 13 | $m(m(x,y,z),x,u)=m(u,y,z)$ | [6 → 9] |
| 23 | $m(m(x,y,z),u,v)=m(m(v,u,x),y,z)$ | [11 → 13] |
| 241 | $m(x,y,m(z,u,v))=m(m(x,y,z),u,v)$ | [6 → 23] |

## Full (gL) vs. Basic (gL)

Here we give an example of a theorem in full (gL) that is not a theorem in basic (gL).

**Theorem.**

$$\left\{\begin{array}{l} x+e=x \\ e+x=x \\ x*(y*x)=y \\ x*y=y*x \end{array}\right\} \;=\!(gL)\!\Rightarrow\; \{x+y=e*(x*y)\}.$$

*Proof* (found by Otter 3.0.5b on soot.mcs.anl.gov at 0.82 seconds).

| | | |
|---|---|---|
| 2 | $x+e=x$ | |
| 3 | $e+x=x$ | |
| 4 | $x*(y*x)=y$ | |
| 5 | $x*y=y*x$ | |
| 11 | $(e+x)*(y*x)=y$ | [3 → 4] |
| 17 | $x*(e+(y*x))=y$ | [3 → 4] |
| 28,27 | $e+(x*y)=y*x$ | [3 → 5] |
| 29 | $x*(x*y)=y$ | [17 :28] |
| 35 | $(x+y)*(z*y)=(x+u)*(z*u)$ | [11 → 11 :(gL)] |
| 53 | $x*(y*e)=(x+z)*(y*z)$ | [2 → 35] |

| | | |
|---|---|---|
| 106 | $(x+y)*((x+z)*y)=z$ | $[35 \to 29]$ |
| 180,179 | $(x+y)*(z*y)=x*(e*z)$ | $[5 \to 53, \text{flip}]$ |
| 194 | $x*(e*(x+y))=y$ | $[106 : 180]$ |
| 225 | $e*(x+y)=x*y$ | $[194 \to 29, \text{flip}]$ |
| 231 | $x+y=e*(x*y)$ | $[225 \to 29, \text{flip}]$ |

To show that the corresponding statement is not a theorem in basic (gL) we used MACE [5], a program that looks for small models or counterexamples of first-order statements. The statements

$$\left\{ \begin{array}{l} (x*y)*z=(x*u)*v \;\to\; (w*y)*z=(w*u)*v \\ x+e=x \\ e+x=x \\ x*(y*x)=y \\ x*y=y*x \\ A+B \neq e*(A*B) \end{array} \right\}$$

have the following model (found by MACE 1.3.2 on ember.mcs.anl.gov at 5.97 seconds).

| $*$ | 0 | 1 | 2 |
|---|---|---|---|
| 0 | 0 | 2 | 1 |
| 1 | 2 | 1 | 0 |
| 2 | 1 | 0 | 2 |

| $+$ | 0 | 1 | 2 |
|---|---|---|---|
| 0 | 0 | 1 | 2 |
| 1 | 1 | 0 | 0 |
| 2 | 2 | 0 | 0 |

e: 0
A: 1
B: 1

# References


[1] D. Eisenbud, M. Green, and J. Harris. Cayley-Bacharach theorems and conjectures. *Bull Amer. Math. Soc (N.S.)*, 33:295–324, 1996.

[2] I. M. S. Etherington. Quasigroups and cubic curves. *Proc. Edinburgh Math. Soc.*, 14:273–291, 1965.

[3] P. Griffiths and J. Harris. *Principles of Algebraic Geometry*. Wiley, New York, 1978.

[4] A. Knapp, editor. *Elliptic Curves*. Princeton University Press, 1993.

[5] W. McCune. Models And Counter-Examples (MACE). http://www.mcs.anl.gov/home/mccune/ar/mace/, 1994.

[6] W. McCune. Otter 3.0 Reference Manual and Guide. Tech. Report ANL-94/6, Argonne National Laboratory, Argonne, IL, 1994.

[7] W. McCune and R. Padmanabhan. *Automated Deduction in Equational Logic and Cubic Curves*, volume 1095 of *Lecture Notes in Computer Science (AI subseries)*. Springer-Verlag, Berlin, 1996.

[8] N. S. Mendelsohn, R. Padmanabhan, and B. Wolk. Straight edge constructions on cubic curves. *C. R. Math. Rep. Acad. Sci. Canada*, 10:77–82, 1988.

[9] J. S. Milne. Abelian varieties. In *Arithmetic, Geometry*, pages 103–150. Springer-Verlag, Berlin, 1986.

[10] D. Mumford. *Abelian Varieties*. Oxford University Press, 1985. Tata Institute of Fundamental Research, Bombay.

[11] R. Padmanabhan. Logic of equality in geometry. *Discrete Mathematics*, 15:319–331, 1982.

[12] R. Padmanabhan and W. McCune. Automated reasoning about cubic curves. *Computers and Mathematics with Applications*, 29(2):17–26, 1995.